


FIG. 2





FIG. 3



10575901 01/08/99

FIG. 5





FIG. 6








FIG. 7

FIG. 8





105193 5120880





FIG. 11
Scfvs to soluble BLYS only






FIG. 12




FIG. 13

FIG. 14
Plate 1079 Sensorgram - 8 Clones





FIG. 15

I082C03 Sensorgram



state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the antibody or fragment thereof, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

[0542] In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it

can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0543] The compositions of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

[0544] The amount of the composition of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, *in vitro* assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0545] For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life within the human body than antibodies from other species due to the immune response to the foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of therapeutic or pharmaceutical compositions of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

[0546] The antibodies and antibody compositions of the invention may be administered alone or in combination with other adjuvants. Adjuvants that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.),

B streptococci. In another embodiment, antibody and antibody compositions of the invention are used in combination with PNEUMOVAX-23™ to treat, prevent, and/or diagnose infection and/or any disease, disorder, and/or condition associated with *Streptococcus pneumoniae*.

[0548] The antibody and antibody compositions of the invention may be administered alone or in combination with other therapeutic agents, including but not limited to, chemotherapeutic agents, antibiotics, antivirals, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents and cytokines. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

[0549] In one embodiment, the antibody and antibody compositions of the invention are administered in combination with other members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), TRAIL, AIM-II (International Publication No. WO 97/34911), APRIL (J. Exp. Med. 188(6):1185-1190 (1998)), endokine-alpha (International Publication No. WO 98/07880), Neutrokine-alpha (International Application Publication No. WO 98/18921), OPG, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

[0550] In a preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with CD40 ligand (CD40L), a soluble form of CD40L (e.g., AVREND™), biologically active fragments, variants, or derivatives of CD40L, anti-CD40L antibodies (e.g., agonistic or antagonistic antibodies), and/or anti-CD40 antibodies (e.g., agonistic or antagonistic antibodies).

[0551] In an additional embodiment, the antibody and antibody compositions of the invention are administered alone or in combination with an anti-angiogenic agent(s). Anti-angiogenic agents that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, Angiostatin (Entremed, Rockville, MD), Troponin-1 (Boston Life Sciences, Boston, MA), anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel (Taxol), Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, VEGI, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

[0552] Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

[0553] Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

[0554] Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include

molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for example, molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

[0555] A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include, but are not limited to, platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., *Cancer Res.* 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydropoline, Thiaproline, alpha, alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., *J. Bio. Chem.* 267:17321-17326, 1992); Chymostatin (Tomkinson et al., *Biochem J.* 286:475-480, 1992); Cyclodextrin Tetradecasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., *Nature* 348:555-557, 1990); Gold Sodium Thiomalate ("GST"; Matsubara and Ziff, *J. Clin. Invest.* 79:1440-1446, 1987); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., *J. Biol. Chem.* 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4- chloroanthronilic acid disodium or "CCA"; (Takeuchi et al., *Agents Actions* 36:312-316, 1992); and metalloproteinase inhibitors such as BB94.

[0556] Additional anti-angiogenic factors that may also be utilized within the context of the present invention include Thalidomide, (Celgene, Warren, NJ); Angiostatic steroid; AGM-1470 (H. Brem and J. Folkman *J Pediatr. Surg.* 28:445-51 (1993)); an integrin alpha v beta 3 antagonist (C. Storgard et al., *J Clin. Invest.* 103:47-54 (1999)); carboxynaminolimidazole; Carboxyamidotriazole (CAI) (National Cancer Institute, Bethesda, MD); Conbretastatin A-4 (CA4P) (OXiGENE, Boston, MA); Squalamine (Magainin Pharmaceuticals, Plymouth Meeting, PA); TNP-470, (Tap Pharmaceuticals, Deerfield, IL); ZD-0101 AstraZeneca (London, UK); APRA (CT2584); Benefin, Byrostatin-1 (SC339555); CGP-41251 (PKC 412); CM101; Dextrazoxane (ICRF187); DMXAA; Endostatin; Flavopridiol; Genestein; GTE; ImmTher; Iressa (ZD1839);

Octreotide (Somatostatin); Panretin; Penacillamine; Photopoint; PI-88; Prinomastat (AG-3340) Purytin; Suradista (FCE26644); Tamoxifen (Nolvadex); Tazarotene; Tetrathiomolybdate; Xeloda (Capecitabine); and 5-Fluorouracil.

[0557] Anti-angiogenic agents that may be administered in combination with the compounds of the invention may work through a variety of mechanisms including, but not limited to, inhibiting proteolysis of the extracellular matrix, blocking the function of endothelial cell-extracellular matrix adhesion molecules, by antagonizing the function of angiogenesis inducers such as growth factors, and inhibiting integrin receptors expressed on proliferating endothelial cells. Examples of anti-angiogenic inhibitors that interfere with extracellular matrix proteolysis and which may be administered in combination with the antibody and antibody compositions of the invention include, but are not limited to, AG-3340 (Agouron, La Jolla, CA), BAY-12-9566 (Bayer, West Haven, CT), BMS-275291 (Bristol Myers Squibb, Princeton, NJ), CGS-27032A (Novartis, East Hanover, NJ), Marimastat (British Biotech, Oxford, UK), and Metastat (Aeterna, St-Foy, Quebec). Examples of anti-angiogenic inhibitors that act by blocking the function of endothelial cell-extracellular matrix adhesion molecules and which may be administered in combination with the antibody and antibody compositions of the invention include, but are not limited to, EMD-121974 (Merck KGaA Darmstadt, Germany) and Vitaxin (Ixsys, La Jolla, CA/Medimmune, Gaithersburg, MD). Examples of anti-angiogenic agents that act by directly antagonizing or inhibiting angiogenesis inducers and which may be administered in combination with the antibody and antibody compositions of the invention include, but are not limited to, Angiozyme (Ribozyme, Boulder, CO), Anti-VEGF antibody (Genentech, S. San Francisco, CA), PTK-787/ZK-225846 (Novartis, Basel, Switzerland), SU-101 (Sugen, S. San Francisco, CA), SU-5416 (Sugen/ Pharmacia Upjohn, Bridgewater, NJ), and SU-6668 (Sugen). Other anti-angiogenic agents act to indirectly inhibit angiogenesis. Examples of indirect inhibitors of angiogenesis which may be administered in combination with the antibody and antibody compositions of the invention include, but are not limited to, IM-862 (Cytran, Kirkland, WA), Interferon-alpha, IL-12 (Roche, Nutley, NJ), and Pentosan polysulfate (Georgetown University, Washington, DC).

[0558] In particular embodiments, the use of antibody and antibody compositions of the invention in combination with anti-angiogenic agents is contemplated for the

treatment, prevention, and/or amelioration of an autoimmune disease, such as for example, an autoimmune disease described herein.

[0559] In a particular embodiment, the use of antibody and antibody compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of arthritis. In a more particular embodiment, the use of antibody and antibody compositions of the invention in combination with anti-angiogenic agents is contemplated for the treatment, prevention, and/or amelioration of rheumatoid arthritis.

[0560] In another embodiment, antibody and antibody compositions of the invention are administered in combination with an anticoagulant. Anticoagulants that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, heparin, warfarin, and aspirin. In a specific embodiment, antibody and antibody compositions of the invention are administered in combination with heparin and/or warfarin. In another specific embodiment, antibody and antibody compositions of the invention are administered in combination with warfarin. In another specific embodiment, antibody and antibody compositions of the invention are administered in combination with warfarin and aspirin. In another specific embodiment, antibody and antibody compositions of the invention are administered in combination with heparin. In another specific embodiment, antibody and antibody compositions of the invention are administered in combination with heparin and aspirin.

[0561] In another embodiment, antibody and antibody compositions of the invention are administered in combination with an agent that suppresses the production of anticardiolipin antibodies. In specific embodiments, the polynucleotides of the invention are administered in combination with an agent that blocks and/or reduces the ability of anticardiolipin antibodies to bind phospholipid-binding plasma protein beta 2-glycoprotein I (b2GPI).

[0562] In certain embodiments, antibody and antibody compositions of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors. Nucleoside reverse transcriptase inhibitors that may be administered in combination with the antibody and antibody compositions of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™

(zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the antibody and antibody compositions of the invention, include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the antibody and antibody compositions of the invention, include, but are not limited to, CRIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with antibody and antibody compositions of the invention to treat, prevent, and/or diagnose AIDS and/or to treat, prevent, and/or diagnose HIV infection.

[0563] In other embodiments, antibody and antibody compositions of the invention may be administered in combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the antibody and antibody compositions of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™, GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ (sargramostim/GM-CSF). In a specific embodiment, antibody and antibody compositions of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat, prevent, and/or diagnose an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat, prevent, and/or diagnose an opportunistic *Mycobacterium avium* complex infection. In another specific embodiment, antibody and antibody compositions of the invention are

used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat, prevent, and/or diagnose an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat, prevent, and/or diagnose an opportunistic cytomegalovirus infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or KETOCONAZOLE™ to prophylactically treat, prevent, and/or diagnose an opportunistic fungal infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat, prevent, and/or diagnose an opportunistic herpes simplex virus type I and/or type II infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat, prevent, and/or diagnose an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, antibody and antibody compositions of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat, prevent, and/or diagnose an opportunistic bacterial infection.

[0564] In a further embodiment, the antibody and antibody compositions of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

[0565] In a further embodiment, the antibody and antibody compositions of the invention are administered in combination with an antibiotic agent. Antibiotic agents that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, amoxicillin, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin, erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin, sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamethoxazole, and vancomycin.

[0566] Conventional nonspecific immunosuppressive agents, that may be

administered in combination with the antibody and antibody compositions of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs cyclophosphamide, cyclophosphamide IV, methylprednisolone, prednisolone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by suppressing the function of responding T cells.

[0567] In specific embodiments, antibody and antibody compositions of the invention are administered in combination with immunosuppressants.

Immunosuppressants preparations that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMUNE™/NEORAL™/SANGDYA™ (cyclosporin), PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucocorticosteroids, and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be used to prevent rejection of organ or bone marrow transplantation.

[0568] In a preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with steroid therapy. Steroids that may be administered in combination with the antibody and antibody compositions of the invention, include, but are not limited to, oral corticosteroids, prednisone, and methylprednisolone (e.g., IV methylprednisolone). In a specific embodiment, antibody and antibody compositions of the invention are administered in combination with prednisone. In a further specific embodiment, the antibody and antibody compositions of the invention are administered in combination with prednisone and an immunosuppressive agent. Immunosuppressive agents that may be administered with the antibody and antibody compositions of the invention and prednisone are those described herein, and include, but are not limited to, azathioprine, cyclophosphamide, and cyclophosphamide IV. In another specific embodiment, antibody and antibody compositions of the invention are administered in combination with methylprednisolone. In a further specific embodiment, the antibody and antibody compositions of the invention are administered in combination with methylprednisolone and an immunosuppressive agent. Immunosuppressive agents that may be administered with the antibody and antibody compositions of the invention and methylprednisolone are those described herein, and include, but are not limited to, azathioprine, cyclophosphamide, and cyclophosphamide IV.

[0569] In a preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with an antimalarial. Antimalarials that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, hydroxychloroquine, chloroquine, and/or quinacrine.

[0570] In a preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with an NSAID.

[0571] In a nonexclusive embodiment, the antibody and antibody compositions of the invention are administered in combination with one, two, three, four, five, ten, or more of the following drugs: NRD-101 (Hoechst Marion Roussel), diclofenac (Dimethaid), oxaprozin potassium (Monsanto), mecamermin (Chiron), T-614 (Toyama), pemetrexed disodium (Eli Lilly), atreleuton (Abbott), valdecoxib (Monsanto), eltenac (Byk Gulden), campath, AGM-1470 (Takeda), CDP-571 (Celltech Chiroscience), CM-101 (CarboMed), ML-3000 (Merckle), CB-2431 (KS Biomedix), CBF-BS2 (KS Biomedix), IL-1Ra gene therapy (Valentis), JTE-522 (Japan Tobacco), paclitaxel (Angiotech), DW-166HC (Dong Wha), darbufelone mesylate (Warner-Lambert), soluble TNF receptor 1 (synergen; Amgen), IPR-6001 (Institute for Pharmaceutical Research), trocade (Hoffman-La Roche), EF-5 (Scotia Pharmaceuticals), BIIL-284 (Boehringer Ingelheim), BIIF-1149 (Boehringer Ingelheim), LeukoVax (Inflammatics), MK-663 (Merck), ST-1482 (Sigma-Tau), and butixocort propionate (WarnerLambert).

[0572] In a preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with one, two, three, four, five or more of the following drugs: methotrexate, sulfasalazine, sodium aurothiomalate, auranofin, cyclosporine, penicillamine, azathioprine, an antimalarial drug (e.g., as described herein), cyclophosphamide, chlorambucil, gold, ENBREL™ (Etanercept), anti-TNF antibody, LJP 394 (La Jolla Pharmaceutical Company, San Diego, California) and prednisolone.

[0573] In a more preferred embodiment, the antibody and antibody compositions of the invention are administered in combination with an antimalarial, methotrexate, anti-TNF antibody, ENBREL™ and/or suflasalazine. In one embodiment, the antibody and antibody compositions of the invention are administered in combination with methotrexate. In another embodiment, the antibody and antibody compositions of the invention are administered in combination with anti-TNF antibody. In another embodiment, the antibody and antibody compositions of the invention are administered in

combination with methotrexate and anti-TNF antibody. In another embodiment, the antibody and antibody compositions of the invention are administered in combination with sufasalazine. In another specific embodiment, the antibody and antibody compositions of the invention are administered in combination with methotrexate, anti-TNF antibody, and sufasalazine. In another embodiment, the antibody and antibody compositions of the invention are administered in combination ENBREL™. In another embodiment, the antibody and antibody compositions of the invention are administered in combination with ENBREL™ and methotrexate. In another embodiment, the antibody and antibody compositions of the invention are administered in combination with ENBREL™, methotrexate and sufasalazine. In another embodiment, the antibody and antibody compositions of the invention are administered in combination with ENBREL™, methotrexate and sufasalazine. In other embodiments, one or more antimalarials is combined with one of the above-recited combinations. In a specific embodiment, the antibody and antibody compositions of the invention are administered in combination with an antimalarial (e.g., hydroxychloroquine), ENBREL™, methotrexate and sufasalazine. In another specific embodiment, the antibody and antibody compositions of the invention are administered in combination with an antimalarial (e.g., hydroxychloroquine), sufasalazine, anti-TNF antibody, and methotrexate.

[0574] In an additional embodiment, antibody and antibody compositions of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the antibody and antibody compositions of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, GAMMAGARD S/D™, and GAMIMUNE™. In a specific embodiment, antibody and antibody compositions of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

[0575] CD40 ligand (CD40L), a soluble form of CD40L (e.g., AVREND™), biologically active fragments, variants, or derivatives of CD40L, anti-CD40L antibodies (e.g., agonistic or antagonistic antibodies), and/or anti-CD40 antibodies (e.g., agonistic or antagonistic antibodies).

[0576] In an additional embodiment, the antibody and antibody compositions of the invention are administered alone or in combination with an anti-inflammatory agent.

Anti-inflammatory agents that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

[0577] In another embodiment, compositions of the invention are administered in combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., mephallen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

[0578] In a specific embodiment, antibody and antibody compositions of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, antibody and antibody compositions of the invention are administered in combination with Rituximab. In a further embodiment, antibody and antibody compositions of the invention are administered with Rituxmab and CHOP, or Rituxmab and any combination of the components of CHOP.

[0579] In an additional embodiment, the antibody and antibody compositions of

the invention are administered in combination with cytokines. Cytokines that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, GM-CSF, G-CSF, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-alpha, IFN-beta, IFN-gamma, TNF-alpha, and TNF-beta. In preferred embodiments, antibody and antibody compositions of the invention are administered with BLYS (e.g., amino acids 134-285 of SEQ ID NO:3228). In another embodiment, antibody and antibody compositions of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, IL-21, and IL-22. In preferred embodiments, the antibody and antibody compositions of the invention are administered in combination with IL4 and IL10.

[0580] In one embodiment, the antibody and antibody compositions of the invention are administered in combination with one or more chemokines. In specific embodiments, the antibody and antibody compositions of the invention are administered in combination with an α (CxC) chemokine selected from the group consisting of gamma-interferon inducible protein-10 (γ IP-10), interleukin-8 (IL-8), platelet factor-4 (PF4), neutrophil activating protein (NAP-2), GRO- α , GRO- β , GRO- γ , neutrophil-activating peptide (ENA-78), granulocyte chemoattractant protein-2 (GCP-2), and stromal cell-derived factor-1 (SDF-1, or pre-B cell stimulatory factor (PBSF)); and/or a β (CC) chemokine selected from the group consisting of: RANTES (regulated on activation, normal T expressed and secreted), macrophage inflammatory protein-1 alpha (MIP-1 α), macrophage inflammatory protein-1 beta (MIP-1 β), monocyte chemotactic protein-1 (MCP-1), monocyte chemotactic protein-2 (MCP-2), monocyte chemotactic protein-3 (MCP-3), monocyte chemotactic protein-4 (MCP-4) macrophage inflammatory protein-1 gamma (MIP-1 γ), macrophage inflammatory protein-3 alpha (MIP-3 α), macrophage inflammatory protein-3 beta (MIP-3 β), macrophage inflammatory protein-4 (MIP-4/DC-CK-1/PARC), eotaxin, Exodus, and I-309; and/or the γ (C) chemokine, lymphotactin.

[0581] In another embodiment, the antibody and antibody compositions of the invention are administered with chemokine beta-8, chemokine beta-1, and/or macrophage inflammatory protein-4. In a preferred embodiment, the antibody and antibody compositions of the invention are administered with chemokine beta-8.

[0582] In an additional embodiment, the antibody and antibody compositions of

the invention are administered in combination with an IL-4 antagonist. IL-4 antagonists that may be administered with the antibody and antibody compositions of the invention include, but are not limited to: soluble IL-4 receptor polypeptides, multimeric forms of soluble IL-4 receptor polypeptides; anti-IL-4 receptor antibodies that bind the IL-4 receptor without transducing the biological signal elicited by IL-4, anti-IL-4 antibodies that block binding of IL-4 to one or more IL-4 receptors, and muteins of IL-4 that bind IL-4 receptors but do not transduce the biological signal elicited by IL-4. Preferably, the antibodies employed according to this method are monoclonal antibodies (including antibody fragments, such as, for example, those described herein).

[0583] The invention also encompasses combining the polynucleotides and/or polypeptides of the invention (and/or agonists or antagonists thereof) with other proposed or conventional hematopoietic therapies. Thus, for example, the polynucleotides and/or polypeptides of the invention (and/or agonists or antagonists thereof) can be combined with compounds that singly exhibit erythropoietic stimulatory effects, such as erythropoietin, testosterone, progenitor cell stimulators, insulin-like growth factor, prostaglandins, serotonin, cyclic AMP, prolactin, and triiodothyronine. Also encompassed are combinations of the antibody and antibody compositions of the invention with compounds generally used to treat aplastic anemia, such as, for example, methenolene, stanozolol, and nandrolone; to treat iron-deficiency anemia, such as, for example, iron preparations; to treat malignant anemia, such as, for example, vitamin B₁₂ and/or folic acid; and to treat hemolytic anemia, such as, for example, adrenocortical steroids, e.g., corticoids. See e.g., Resegotti et al., *Panminerva Medica*, 23:243-248 (1981); Kurtz, *FEBS Letters*, 14a:105-108 (1982); McGonigle et al., *Kidney Int.*, 25:437-444 (1984); and Pavlovic-Kantera, *Expt. Hematol.*, 8(supp. 8) 283-291 (1980), the contents of each of which are hereby incorporated by reference in their entireties.

[0584] Compounds that enhance the effects of or synergize with erythropoietin are also useful as adjuvants herein, and include but are not limited to, adrenergic agonists, thyroid hormones, androgens, hepatic erythropoietic factors, erythrotropins, and erythroginins. See for e.g., Dunn, "Current Concepts in Erythropoiesis", John Wiley and Sons (Chichester, England, 1983); Kalmani, *Kidney Int.*, 22:383-391 (1982); Shahidi, *New Eng. J. Med.*, 289:72-80 (1973); Urabe et al., *J. Exp. Med.*, 149:1314-1325 (1979); Billat et al., *Expt. Hematol.*, 10:133-140 (1982); Naughton et al., *Acta Haemat.*, 69:171-

179 (1983); Cognote et al. in abstract 364, Proceedings 7th Intl. Cong. of Endocrinology (Quebec City, Quebec, July 1-7, 1984); and Rothman et al., 1982, J. Surg. Oncol., 20:105-108 (1982). Methods for stimulating hematopoiesis comprise administering a hematopoietically effective amount (i.e., an amount which effects the formation of blood cells) of a pharmaceutical composition containing polynucleotides and/or polypeptides of the invention (and/or agonists or antagonists thereof) to a patient. The polynucleotides and/or polypeptides of the invention and/or agonists or antagonists thereof is administered to the patient by any suitable technique, including but not limited to, parenteral, sublingual, topical, intrapulmonary and intranasal, and those techniques further discussed herein. The pharmaceutical composition optionally contains one or more members of the group consisting of erythropoietin, testosterone, progenitor cell stimulators, insulin-like growth factor, prostaglandins, serotonin, cyclic AMP, prolactin, triiodothyronine, methenolone, stanozolol, and nandrolone, iron preparations, vitamin B₁₂, folic acid and/or adrenocortical steroids.

[0585] In an additional embodiment, the antibody and antibody compositions of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

[0586] In an additional embodiment, the antibody and antibody compositions of the invention are administered in combination with fibroblast growth factors. Fibroblast growth factors that may be administered with the antibody and antibody compositions of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

[0587] Additionally, the antibody and antibody compositions of the invention may be administered alone or in combination with other therapeutic regimens, including but not limited to, radiation therapy. Such combinatorial therapy may be administered sequentially and/or concomitantly.

Kits

[0588] The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the pharmaceutical

compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

[0589] The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In an alternative embodiment, a kit comprises an antibody fragment that immunospecifically binds to BLyS. In a specific embodiment, the kits of the present invention contain a substantially isolated BLyS polypeptide as a control. Preferably, the kits of the present invention further comprise a control antibody which does not react with BLyS. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to BLyS (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes the first antibody may be conjugated to a detectable substrate). In specific embodiments, the kit may include a recombinantly produced or chemically synthesized BLyS. The BLyS provided in the kit may also be attached to a solid support. In a more specific embodiment the detecting means of the above-described kit includes a solid support to which BLyS is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of the antibody to BLyS can be detected by binding of the said reporter-labeled antibody.

[0590] In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with BLyS, and means for detecting the binding of BLyS to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

[0591] In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound BLyS obtained by the methods of the present invention. After BLyS binds to a specific antibody, the unbound serum components are removed by

washing, reporter-labeled anti-human antibody is added, unbound anti-human antibody is removed by washing, and a reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-BLyS antibody on the solid support. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate.

[0592] The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

[0593] Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant BLyS, and a reporter-labeled anti-human antibody for detecting surface-bound anti-BLyS antibody.

[0594] In specific embodiments, the present invention encompasses a single chain Fv (scFv) having an amino acid sequence of one of SEQ ID NOS: 1 to 2128.

[0595] In specific embodiments, the present invention encompasses a single chain Fv (scFv) having an amino acid sequence of one of SEQ ID NOS: 1 to 46, 321 to 329, 1563 to 1595, and 1881 to 1908.

[0596] In specific embodiments, the present invention encompasses a single chain Fv (scFv) having an amino acid sequence of one of SEQ ID NOS: 1563 to 1880.

[0597] In specific embodiments, the present invention encompasses a single chain Fv (scFv) having an amino acid sequence of one of SEQ ID NOS: 1881 to 2128.

[0598] In specific embodiments, the present invention encompasses a single chain Fv (scFv) having an amino acid sequence of one of SEQ ID NOS: 1 to 1562.

[0599] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VH domain from an scFv having an amino acid sequence

of one of SEQ ID NOS: 1 to 2128, wherein said antibody or fragment thereof immunospecifically binds BLyS.

[0600] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VH domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1 to 46, 321 to 329, 1563 to 1595, and 1881 to 1908.

[0601] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VH domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1881 to 2128, and in which said antibody or fragment thereof immunospecifically binds to the membrane-bound form of BLyS.

[0602] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VH domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1563 to 1880, and in which said antibody or fragment thereof immunospecifically binds to the soluble form of BLyS.

[0603] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VL domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1 to 2128, wherein said antibody or fragment thereof immunospecifically binds BLyS.

[0604] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VL domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1 to 46, 321 to 329, 1563 to 1595, and 1881 to 1908.

[0605] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VL domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1881 to 2128, and in which said antibody or fragment thereof immunospecifically binds to the membrane-bound form of BLyS.

[0606] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VL domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1563 to 1880, and in which said antibody or fragment thereof immunospecifically binds to the soluble form of BLyS.

[0607] In specific embodiments, the present invention encompasses an antibody or fragment thereof comprising a VL domain from an scFv having an amino acid sequence of one of SEQ ID NOS: 1 to 2128, wherein said antibody or fragment thereof immunospecifically binds BLyS and which also comprises a VH domain from an scFv

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I declare that I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

ANTIBODIES THAT IMMUNOSPECIFICALLY BIND TO BLYS

the specification of which is attached hereto or was filed on (June 15, 2001) herewith as Application Serial No. not yet assigned and was amended on _____ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim foreign priority benefits under 35 U.S.C. § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application, which designated at least one country other than the United States listed below, and have also identified below any foreign application for patent or inventor's certificate, or PCT international application having a filing date before that of the application on which priority is claimed. Prior Foreign Application(s):

Priority Claimed
Yes No

(Number)

(Country)

(Day/Month/Year Filed)

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below:

60/212,210

(Provisional App. Serial No.)

June 15, 2000

(Filing Date)

60/240,816

(Provisional App. Serial No.)

October 17, 2000

(Filing Date)

60/276,248

(Provisional App. Serial No.)

March 16, 2001

(Filing Date)

60/277,379

(Provisional App. Serial No.)

March 21, 2001

(Filing Date)

60/293,499

(Provisional App. Serial No.)

May 25, 2001

(Filing Date)

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s) or under § 365(b) of any PCT international application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Section 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, Section 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status: patented, pending,
abandoned)

I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: James H. Davis (Reg. No. 40,582), Kenley K. Hoover (Reg. No. 40,302), Michele M. Wales (Reg. No. 43,975), Jonathan L. Klein (Reg. No. 41,119), Mark J. Hyman (Reg. No. 46,789), and Joseph J. Kenny (Reg. No. 43,710) of Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, Maryland, 20850.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first joint inventor: Steven M. RUBEN

Inventor's signature: _____ Date: _____
Residence: 18528 Heritage Hills Drive, Olney, Maryland 20832 Citizenship: US
Post Office Address: same as above

Full name of additional joint inventor: Steven C. BARASH

Inventor's signature: _____ Date: _____
Residence: 111 Watkins Pond Blvd., #301, Rockville, MD 20850 Citizenship: US
Post Office Address: same as above

Full name of additional joint inventor: Gil H. CHOI

Inventor's signature: _____ Date: _____
Residence: 11429 Potomac Oaks Drive Citizenship: KR
Post Office Address: same as above

Full name of additional joint inventor: Tristan VAUGHAN

c/o Cambridge Antibody Technology Group PLC

The Science Park, Melbourne, Nr Royston, Cambridgeshire SG8 6JJ

Inventor's signature: _____ Date: _____
Residence: UK Citizenship: UK
Post Office Address: same as above

Full name of additional joint inventor: David HILBERT

Inventor's signature: 8501 Meadowlark Lane, Bethesda, MD 20817 Date: US
Residence: _____ Citizenship: _____
Post Office Address: same as above